

Mr Mark S. Falworth MBBS FRCS(Eng) FRCS (Orth)
Consultant Orthopaedic Surgeon
Shoulder and Elbow Surgery

Spire Bushey Hospital
Heathbourne Road
Bushey
Watford
Herts, WD23 1RD

Royal National Orthopaedic Hospital
Brockley Hill
Stanmore
Middlesex
HA7 4LP

The Stanmore Clinic
HCA Elstree Waterfront
The Waterfront
Beaufort House, Elstree Rd.
Herts, WD6 3BS

Appointments & Queries:
Practice Manager: Lynn Etherton
Telephone: 020 7042 1807
Fax: 020 7042 1808
Email: lynn.ertherton@phf.uk.com

All correspondence to Spire Bushey Hospital

Physiotherapy protocol: Revision Total Shoulder Replacement

Weeks 1-6:

- Sling for 6 weeks
- No active IR /hand behind back / backwards extension for 6 weeks (so as to protect the subscapularis repair.)
- ROM goals: Week 1: 90° FE (off the brace in the scapula plane), 0° ER at side; ABD max 75° without rotation
- ROM goals: Week 3: 120° FE (off the brace in the scapula plane), 20° ER at side; ABD max 75° without rotation
- Canes/pulleys OK to advance from PROM
- PROM → AAROM
- Heat before PT, ice after PT

Weeks 6-12:

- Begin AAROM → AROM for internal rotation and backwards extension as tolerated, if not already begun.
- Goals: Increase ROM as tolerated with gentle passive stretching at end ranges
- Begin light resisted ER/FE/ABD: isometrics and bands, concentric motions only
- No resisted internal rotation/backwards extension until 12 weeks post-op
- No scapular retractions with bands yet

Months 3-12:

- Increase resisted IR/ER/FE/ABD (isometrics/bands): isometrics → light bands → weights
- Advance strengthening as tolerated; 10 reps/1 set per exercise for rotator cuff, deltoid, and scapular stabilizers.
- Increase ROM to full with passive stretching at end ranges
- Begin eccentric motions and closed chain exercises at 12 weeks.

Comments: