

Mr Mark S. Falworth MBBS FRCS(Eng) FRCS (Orth)
Consultant Orthopaedic Surgeon
Shoulder and Elbow Surgery

Spire Bushey Hospital
Heathbourne Road
Bushey
Watford
Herts, WD23 1RD

Royal National Orthopaedic Hospital
Brockley Hill
Stanmore
Middlesex
HA7 4LP

The Stanmore Clinic
HCA Elstree Waterfront
The Waterfront
Beaufort House, Elstree Rd.
Herts, WD6 3BS

Appointments & Queries:
Practice Manager: Lynn Etherton
Telephone: 020 7042 1807
Fax: 020 7042 1808
Email: lynn.ertherton@phf.uk.com

All correspondence to Spire Bushey Hospital

Physiotherapy protocol: Rotator Cuff Repair + subacromial decompression

Weeks 0-6:

- Elbow ROM, wrist ROM, grip strengthening and scapular setting exercises
- Patient to remain in sling for 6 weeks
- True PROM only - the rotator cuff tendon needs to heal back into the bone
- ROM goals: 110° FE, 0° ER at side; ABD max 60-80° without rotation
- No resisted motion of shoulder until 12 weeks post-op
- Minimise canes/pulleys until 6 weeks post-op, because these are active-assisted exercises
- Heat before Physio, ice / cryocuff after

Weeks 6-12:

- Begin AAROM as tolerated
- Goals: re-establish full range of movement with 160 FE, 30 ER increasing as tolerated
- Light passive stretching at end ranges
- At 8 weeks, can begin isometric strengthening to unaffected muscle groups

Months 3-12:

- Advance to full AROM as tolerated with passive stretching at end ranges
- Advance strengthening as tolerated: isometrics → bands → light weights (1-5 lbs); 8-12 reps/2-3 sets per rotator cuff, deltoid, and scapular stabilisers
- Be careful not to overdue strengthening to avoid rotator cuff tendonitis (3x/week)
- Begin eccentrically resisted movement and exercises as able

Comments: